

RESEARCH

Open Access



How has the COVID-19 epidemic affected the risk behaviors of people who inject drugs in a city with high harm reduction service coverage in Vietnam? A qualitative investigation

Trang Thu Nguyen^{1*} , Giang Thi Hoang², Duc Quang Nguyen², Anh Huu Nguyen¹, Ngoc Anh Luong¹, Didier Laureillard^{3,4}, Nicolas Nagot³, Don Des Jarlais⁵, Huong Thi Duong², Thanh Thi Tuyet Nham⁶, Oanh Thi Hai Khuat⁶, Khue Minh Pham², Mai Sao Le², Laurent Michel⁷, Delphine Rapoud³ and Giang Minh Le¹

Abstract

Introduction: The COVID-19 outbreak disproportionately affects vulnerable populations including people who inject drugs (PWID). Social distancing and stay-at-home orders might result in a lack of access to medical and social services, poorer mental health, and financial precariousness, and thus, increases in HIV and HCV risk behaviors. This article explores how the HIV/HCV risk behaviors of PWID in Haiphong, a city with high harm reduction service coverage in Vietnam, changed during the early phase of the COVID-19 pandemic, and what shaped such changes, using the risk environment framework.

Method: We conducted three focus group discussions with peer outreach workers in May 2020 at the very end of the first lockdown, and 30 in-depth interviews with PWID between September and October 2020, after the second wave of infection in Vietnam. Discussions and interviews centered on the impact of the COVID-19 pandemic on their lives, and how their drug use and sexual behaviors changed as a result of the pandemic.

Results: The national shutdown of nonessential businesses due to the COVID-19 epidemic caused substantial economic challenges to participants, who mostly were in a precarious financial situation before the start of the epidemic. Unsafe injection is no longer an issue among our sample of PWID in Haiphong thanks to a combination of different factors, including high awareness of injection-related HIV/HCV risk and the availability of methadone treatment. However, group methamphetamine use as a means to cope with the boredom and stress related to COVID-19 was common during the lockdown. Sharing of smoking equipment was a standard practice. Female sex workers, especially those who were active heroin users, suffered most from COVID-related financial pressure and may have engaged in unsafe sex.

Conclusion: While unsafe drug injection might no longer be an issue, group methamphetamine use and unsafe sex were the two most worrisome HIV/HCV risk behaviors of PWID in Haiphong during the social distancing and

*Correspondence: trangthu@hmu.edu.vn

¹ Centre for Training and Research on Substance Use and HIV, Hanoi Medical University, Room 211B, Building E3, #1 Ton That Tung Street, Hanoi, Vietnam

Full list of author information is available at the end of the article



lockdown periods. These elevated risks could continue beyond the enforced lockdown periods, given PWID in general, and PWID who are also sex workers in particular, have been disproportionately affected during the global crisis.

Keywords: COVID-19, Risk behaviors, People who inject drugs, Harm reduction

Introduction

The current outbreak of the novel coronavirus SARS-CoV-2 (COVID-19) has spread all over the world since the first case was detected in December 2019. The pandemic poses a serious threat to global health with significant loss of life and the impact on the society as a whole through the mitigation measures taken to control it. It disproportionately affects vulnerable populations, including people who inject drugs (PWID), of which the majority live in poor socioeconomic conditions [1].

Existing literature points to multiple factors at different levels that may contribute to the harms PWID currently experience during the COVID-19 pandemic [2–6]. People who use drugs are particularly vulnerable to SARS-CoV-2 infection and complications due the high prevalence co-morbidities in their population [5–8]. Many have lung and liver diseases that may dangerously contribute to the aggravation of COVID-related disease [3]. Moreover, the high prevalence of HIV infection in this population indicates that some have weakened immunity and low resistance to COVID infection [3, 7]. A large number of people who use drugs also lack access to personal protection measures such as private space to effectively self-isolate [9]. Job loss or reduced income might cause them great challenges, especially those living hand-to-mouth [1]. Restrictions on face-to-face contact, reduced number of visits per day, or even temporary closures of harm reduction services might result in a lack of timely support for PWID, thus increasing HIV and HCV risk behaviors [3, 10, 11]. Lastly, social isolation and stress might exacerbate negative effects on their mental health [3, 12].

The link between the COVID-19 pandemic and HIV/HCV risk behaviors is little studied. While traditional risk factors like injection equipment sharing and unsafe sexual behaviors might decrease as a result of physical distancing and curfew implementation in many countries [13], these measures might also introduce new risks. A study showed that while sexual risk behaviors among sexual minority men overall reduced during the COVID-19 pandemic, such behaviors increased among those who use drugs [14]. A report from Thailand also suggested increase in sexualized drug use among gay and bisexual men which could lead to sexual risk behaviors [15]. People might also increase their consumption of alcohol and other drugs to cope with the stress related to the pandemic, or get involved with informal economies like

sex work to overcome financial challenges [3, 5, 10]. As COVID-19 public health responses are stretching the already limited resources for the prevention and treatment of drug-related issues, a good understanding of the situation is critical to implement cost-effective harm reduction strategies.

Injection drug use, and more recently unsafe sex, remain the drivers of the HIV/HCV epidemic in Vietnam [16]. Harm reduction programs have been implemented in the country since the mid-1990s, as HIV prevalence among PWID rocketed from 10.1% in 1996 to 32% in 2002 [17]. Major strategies included the provision of sterile needles and syringes, condom distribution and methadone maintenance treatment [18]. Over more than a decade, methadone treatment has expanded from two clinics serving nearly 1000 patients in 2008, to 316 clinics in all provinces, serving about 53,000 (28%) people who use opioids in the country [19]. These programs reduced HIV prevalence among PWID to 12.7% in 2019 [19]. However, while HIV incidence in this group has decreased, their HCV incidence remains as high as 19.4/100 PY [20]. Moreover, increasing consumption of methamphetamine (hereafter: meth) among PWID, including methadone patients in recent years, is raising concerns about a resurgence of HIV/HCV infection and suboptimal treatment outcomes [19, 21–23].

The COVID-19 epidemic started in late January 2020 in Vietnam [24]. Despite the relatively small number of cases (254 by April 10th, 2020), Vietnam conducted the first nationwide lockdown for 3 weeks from April 1st, 2020 as an early response to the epidemic [25]. Non-essential businesses were shut down and medical professionals of different expertise were deployed to scale up COVID-19 prevention and treatment. Since then, subsequent lockdowns for extended periods have continued. Our study explores how the HIV and HCV risk behaviors of PWID changed during the COVID-19 pandemic in Haiphong—a socioeconomically important province in Northern Vietnam with a past drug injection-driven HIV epidemic—and what shaped such changes. At the time of the study, there were no confirmed COVID-19 cases in Haiphong. Nevertheless, the province was under the effect of national measures to contain the COVID-19 pandemic.

Since the early days of the HIV epidemic, the province has implemented various harm reduction strategies [26]. Haiphong is the study site of the 5-year DRIVE (DRug

use and Infections in ViEtnam) project and its associated studies, which have aimed to prevent HIV/HCV transmission among PWID (heroin) in the province since 2016 [27]. This has included using a mass screening strategy coupled with HIV/HCV sensitization, harm reduction, and referral to HIV/HCV, methadone and psychiatric treatments by community-based outreach workers. In total, these studies have reached 4000 out of about 5000 active PWID in the province. Within these projects, 1000 PWID have been initiated on HCV treatment, and about 250 PWID presenting with a mental health diagnosis have received psychiatric treatment. Intensive harm reduction efforts have brought HIV incidence among PWID in Haiphong to a very low level, supporting the declaration of the end of the HIV epidemic among this population in the province [28]. The high number of PWID in HCV treatment also differentiates Haiphong from other provinces in terms of access to HCV care [29]. Given such achievements, Haiphong served as a good study context to understand the early impact of the COVID-19 pandemic on risk behaviors in PWID.

To investigate what influenced risk behaviors, we employed the risk environment framework [30]. This framework has gained popularity in public health over the last 2 decades. Risk environment can be defined as a 'space—whether social or physical—in which a variety of factors interact to increase the chances of drug-related harms' [30]. This framework emphasizes the role of contextual factors in producing or reducing drug-related harms, and provides an alternative to the biomedical and political perspectives that stress individual responsibility in managing harms [31, 32]. The risk environment framework delineates four types of environments—physical, social, political and economic—that interact among themselves and with the individual at micro and macro levels [30]. Identifying both risk and protective environmental factors, and the ways they shape PWID's risk behaviors, is a straightforward way to foster 'enabling environments' for harm reduction [32].

Method

This qualitative analysis is part of DRIVE-COVID, a mixed-methods research project that has investigated the changes in risk behaviors of PWID in Vietnam during the COVID-19 pandemic. We conducted three focus group discussions (FGDs) with peer outreach workers in May 2020 at the end of the first lockdown, and 30 in-depth interviews with PWID between September and October 2020, after the second wave of infection in the country. Peer outreach workers came from the three largest community-based organizations in Haiphong serving people who use drugs and female sex workers. We selected participants to cover a variety of characteristics including

gender, age group, methadone treatment and HIV status, mental health treatment, and HCV treatment status. These characteristics have been reported in systematic reviews to have an association with the HIV/HCV risk behaviors [33–35].

The discussions with peer outreach workers centered on the impact of the COVID-19 pandemic on their lives, and their observation of changes in their clients' injection and sexual risk behaviors. Interviews with participants who use drugs explored their experiences with the pandemic and the lockdown period, with a focus on how their drug use and sexual behaviors changed. The first author developed the FGDs and interview schedules from the research questions with inputs from the whole team. FGD and interviews were conducted in the offices of two community-based organizations that implement peer outreach. All FGDs and interviews were audio-recorded and transcribed verbatim. After being de-identified, transcripts were uploaded onto Atlas.ti version 8 [36] to facilitate coding. Participants received VND200,000 (~US 9 dollars) for their time and effort. The study was approved by the Ethics Committee of Haiphong Medicine and Pharmacy University.

Our analysis was informed by the grounded theory approach. The authors at Hanoi Medical University conducted FGDs and interviews, and summarized salient themes as each session was completed. At first, we read the transcripts multiple times to immerse ourselves in participants' stories. We then conducted open coding until no new codes emerged (after 5 transcripts) [37]. Examples of codes included 'poor people die first', 'COVID-related financial constraints', 'challenging to purchase drugs' or 'care for clients during the lockdown'. We grouped the codes into larger categories of risk and protective environmental factors influencing risk behaviors. The qualitative findings were discussed within the study team and triangulated with the quantitative data on the description of participants' risk behaviors.

Results

Table 1 presents the participants' sociodemographic characteristics. Women accounted for 57% of the sample. All participants self-identified as heterosexual. The median age of participants was 40.5 years. One third of participants were HIV-positive. Two thirds were unemployed or had unstable incomes. 60% were undergoing methadone treatment. Almost half of participants were actively using heroin (daily or once every several days) and one third were actively using meth (once every several days) at the time of the study.

The analysis centered on heroin injection, meth use in groups, and unsafe sexual behaviors as the key HIV/HCV risk behaviors among our participants. Participants

Table 1 Participant characteristics (N=30)

Male/Female	13/17
Transgendered people	0
Sex workers	5
Age (median)	40.5
Education	
Primary school	3
Middle school	10
High school & college	10
No information	7
Occupation	
Unemployed	7
Unstable income	13
Stable income	10
Marital status	
Single	7
Married/Cohabiting	12
Divorced/Widowed	11
Received mental health treatment	15
Received HCV treatment	10
HIV positive/Under antiretroviral treatment	9/9
Currently in methadone treatment	18
Currently using heroin	14
Currently using meth	10

reported no meth injection. The COVID-19 epidemic and containment policies introduced significant changes into the lives of participants. These changes exerted both negative and positive impacts on risk behaviors. Table 2 summarizes the risk and protective environmental factors influencing the three identified behaviors.

Impact of the COVID-19 lockdown on participants’ lives and outreach work

While the lockdown could affect participants in all aspects of life, participants were most worried about its financial impact. All except one participant reported being challenged financially by the COVID-19 outbreak. Because of the lockdown, most socioeconomic activities ceased. Many participants with jobs like waste pickers, waiters, tea stall owners or security guards had to stay at home without work. Some lived hand-to-mouth and struggled to support themselves and their families. Factory workers, despite not losing their jobs, had to significantly reduce their working hours, and with it, their salaries. Few participants who had their ID card and were qualified as being ‘poor’ received some government support. But such support (about five kilograms of rice per person for 3 weeks of lockdown) was insufficient. A participant expressed her desperation:

It’s so stressful. You have no money. You can make no money. You can’t go out to steal; you can’t go out to sell sex. Everyone is dead. (Female, 42 years old, unemployed)

Another person highlighted that poor people were most vulnerable to the impacts of lockdown, noting that social inequality seems to be most pronounced in times of crisis:

For waste pickers, there is no junk to pick in COVID time [...] For people who have money, during these 15, 20 days of lockdown, they could have some rest at home; for those without a decent job, they can make no money for food. (Female, 54 years old, waste picker)

Participants who were active users and were not already under methadone treatment were hit hardest. These people previously had jobs and relatively stable incomes to support their habit. However, as economic activities slowed down, their income also declined. All had to reduce their intake and in some way suffered the discomfort related to insufficient dosage. Participants expressed their anxiety when facing the inability to procure drugs:

When I saw on the news quarantined people [in another country] crying and screaming, I was so upset. In quarantine, money serves nothing. You won’t be able to buy drugs. [Sigh] You’re not addicted, you don’t know ... (Female, 38 years old, sex worker)

Some participants reported that financial challenges caused family conflicts, and both were great stressors to them. Compared to the urgent concern for earning a living, the risk of COVID infection was not a priority. Indeed, many participants reported that they did not worry about contracting COVID while they had to endure hardship in their everyday lives.

Heroin injection

Among the 14 participants who were active heroin users at the time of the interview, 8 were not undergoing methadone treatment. Still, all of these active heroin users reported no sharing of injection equipment. This finding is consistent with our survey data, with 99.5% participants reported not having shared or divided drugs with a used needle or syringe over the last 6 months [38]. The protective social and policy factors against sharing of injection equipment included awareness of the HIV/HCV risks relating to unsafe injection, ease of buying equipment at pharmacies, family support and high methadone treatment coverage. A participant expressed her

Table 2 Summary of risk and protective environmental factors for drug use and sexual behaviors during COVID-19 lockdown

Risk factors	Protective factors
Heroin injection	
<p>Social Lockdown Fear of being arrested by the police when going to buy syringes or drugs</p> <p>Economic Higher price of heroin</p>	<p>Social High awareness of injection-related HIV/HCV risks Ease of buying syringes and needles at pharmacies, even during the lockdown Family support Social norms around heroin Hiding one's drug use status</p> <p>Policy Availability of methadone treatment</p> <p>Physical Police presence in the street, curfew</p> <p>Economic Higher price of heroin</p>
Group methamphetamine use	
<p>Social Social norms related to meth use</p> <p>Physical Having more free time Effects of meth</p> <p>Sexual behaviors</p> <p>Social Fear of COVID-19 infection</p> <p>Physical Curfew, social distancing</p> <p>Economic Reduced income</p>	<p>Social Hiding one's drug use status</p> <p>Economic Reduced income</p> <p>Social High awareness of HIV risk sexual behaviors</p>

concerns about the HCV infection risk between her son and her. She reported how she instructed her son to use drugs safely:

I tell him to use a syringe once and that only when he runs out of money can he use it twice. Each time he would need to wash it with clean water, put it in a safe place and never share it with anyone. My HCV infection has been cured, his hasn't. (Female, 54 years old, waste picker)

Even during the lockdown, participants were still able to buy sterile syringes at pharmacies. A slight increase in price in some pharmacies did not interfere with their practice. Participants often bought syringes for each day's worth of use. Some family even assumed this task and bought boxes of syringes for participants so that they did not have to go out during the lockdown.

The safe injection practice could also have been fostered by the fact that some participants insisted on using heroin alone, since they did not want to disclose their heroin use to their drug using peers:

I've been doing heroin alone since I relapsed. I don't want my friends to know about it. They know I'm doing meth but for them, heroin users are worse than meth users. (Male, 37 years old, unemployed)

Doing heroin alone, however, could increase the risk of overdose death [39].

During lockdown, some participants had to reduce their heroin intake because of their income reduction and the challenges in purchasing heroin. Some did not want their family members to know they were still using heroin. Participants undergoing methadone treatment encountered no difficulty in temporarily stopping or reducing their heroin consumption, while those not undergoing treatment struggled. One participant described the difficulties in buying drugs:

During the lockdown, the police were everywhere. Neither we nor the dealers were allowed to go out. It was too easy to get caught if you went to buy drugs. (Male, 27 years old, painter)

These challenges made him decide to get methadone treatment:

I've known about methadone programs for two years but I didn't really pay attention to it. But given the last lockdown period, I made up my mind and got treatment. (Male, 27 years old, painter)

Outreach workers were, however, less confident that everyone could practice safe injection during the social distancing period and lockdown. They explained:

The dealers no longer sold drugs in small quantities. Several people had to pool their money to buy drugs together. However, since sharing dry drugs results in unequal partitioning, they'd mix drugs with water in a new syringe. Some without money would have to use others' needles. (FGD with peer outreach workers)

Thus, from the outreach workers' perspective, the same financial challenges in buying drugs could facilitate unsafe injection.

Other participants also mentioned the prospect of being arrested by the police while going out to buy sterile syringes at pharmacies.

I know there are many cops around the pharmacy at Hospital X. They'd stay behind you. When they'd see you were buying syringes, they'd get you. That's why I've never bought [syringes] there. (Female, 23 years old, assistant mason)

While there was no report of its impact on injection behaviors, this fear might play a role in the fact that some participants reused their own syringes to avoid going out too often, and thus increased their risk of bacterial infection [40].

Group meth use

Most participants had experience using meth. Among them, 10 people were active meth users and 2 were HCV-positive. Participants often used meth in social gatherings and shared the same smoking tools (bongs and pipes). Only two participants reported doing meth exclusively by themselves. The tools were not disinfected after each use and were replaced every 1 or 2 months. The social and physical factors facilitating meth use in groups included social norms related to meth, having more free time, and the effects of meth. Most participants used meth to enhance friendly conversations.

There are always two, three people or more smoking ice together. Only meth addicts do it by themselves. People normally gather to talk. (Female, 42 years old, sex worker)

Typically, individuals did not just pay for their own consumption but took turns to purchase the drugs for each other. Some stated that they had never had to pay since they only did meth when their friends asked them to join.

The lockdown and social distancing did not seem to reduce meth use. Most participants reported little changes in their patterns of meth use. Having more free time during the lockdown meant more meth-smoking gatherings for some people.

[Before social distancing,] I was busy with my work. I didn't think about [meth] or I wasn't at home to join my friends. During the lockdown, we could go nowhere and had nothing to do. We called each other and went to a friend's house. (Male, 27 years old, painter)

Outreach workers also observed increases in meth use among female sex workers:

In some places, before there was only one ice smoker, but now everyone smokes it. [...] People are more indebted to their patrons [because the patrons bought them drugs and food during the lockdown]. (FGD with peer outreach workers)

Five participants indicated the euphoric effects of meth and another reported meth-related hallucinations as physical risk factors that made them neglect the HCV, COVID-19 and other respiratory infection risks associated with these gatherings. When being asked how they perceived the risk related to sharing smoking pipes, some participants reported they did not even think about it:

My mind was on so many other things then. I was even hallucinating. (Male, 43 years old, unemployed)

Others said they were somewhat afraid of being infected with COVID-19 but they believed that they and their friends could not be infected:

Of course I'm afraid of COVID. But I trust them. (Female, 33 years old, sex workers)

On the other hand, the main protective factors against group meth use were COVID-related financial difficulties and their desire to hide their drug use from the family:

I used to do meth but now I don't, because I've no money and if I do meth now, my parents will know and I don't like it. I force myself to stay at home. (Female, 35 years old, unemployed)

Sexual behaviors

Most male participants reported having lost interest in sex after years of drug use or as a side effect of methadone treatment. Compared to them, women were much more sexually active. Among the five women who reported selling sex occasionally or professionally, one had accepted unsafe sex in exchange for money during the lockdown. Risk factors were related to the COVID-19 epidemic.

Being afraid of COVID-19 infection during intimate transactions, many clients ceased visiting sex workers. The few clients who continued these visits paid sex workers less for the reason that clients earned less too. The curfew order requiring people to stay at home after 10 p.m. directly interfered with sex work since sexual transactions mostly occurred at nighttime. The resulting income loss made sex workers, especially those who were active heroin users, vulnerable, since they could no longer select their clients under the pressure of earning money for the next fix. One participant reported that she had to accept clients she deemed at-risk and had unsafe sex with them when they requested. The prospect of being in withdrawal was terrifying to her. She described her anxiety:

Only seeing them get into the room made me feel reassured. If they shook their head, I'd be nervous. Without a client, I'll be dead. (Female, 38 years old, sex worker)

One protective factor was participants' awareness of the HIV risk related to unsafe sex. In non-COVID times, most of these women insisted on consistent condom use with their clients. One woman accepted unprotected sex with her long-term elder clients but only after she had considered all potential risks. A participant said she almost hit a client when she found out he removed his condom during intercourse:

Who knows how many sex workers he's gone with?! Around here I know three gals with AIDS. [...] One or two hundred [thousand Dong] is not worth getting the disease. (Female, 33 years old, sex worker)

Thus, while participants were well aware of the HIV risk, financial challenges and the urge for heroin consumption might have forced them to resort to unsafe sex.

Discussion

While PWID have been disproportionately affected by the COVID-19 epidemic, little is known about how their HIV/HCV risk behaviors change as a result of the epidemic, and the measures taken to contain it. This study is among the first studies exploring this topic in Vietnam.

It highlights both risk and protective environmental factors influencing the risk behaviors of PWID in the unique context of Haiphong, a province with high harm reduction service coverage in the early phase of the COVID-19 epidemic in Vietnam. The findings herein are critical for harm reduction programs to support PWID during this global crisis.

The challenges that our participants experienced echo what has been found in other studies [1, 2]. Globally, PWID belong to the populations of low socioeconomic status who have been most disadvantaged during the COVID-19 crisis [1]. In Vietnam, as adult children, especially sons, cohabit with their parents to take care of them, most of our participants were living with their family during the lockdown. Thus, they could be more materially and emotionally supported than PWID in other parts of the world, who experienced homelessness and inadequate housing, making them vulnerable to COVID-19 infection and other life hazards [41, 42]. Nonetheless, Vietnamese PWID struggled financially to support themselves and their families. Since the social networks of PWID often consist of people of similarly low socioeconomic status [43], their usual support resources were also depleted under the impact of COVID-19. The epidemic made the experiences of social inequality more pronounced, as our participants observed that the middle class, with savings and the ability to work from home, are better protected from COVID-related financial crises than blue-collar workers. This resonates with Nanda's writing about inequalities during the COVID-19 outbreak [1].

Unsafe injection did not seem to be an issue among PWID in Haiphong, thanks to the high awareness of participants regarding injection-related HIV/HCV risk. However, this might not be the case in provinces with lower harm reduction service coverage. Haiphong PWID did not seem to encounter challenges in accessing harm reduction services during the pandemic, unlike PWID in some European countries [3]. Pharmacies remained open during the lockdown, providing easy access to injection equipment. However, the punitive approach of the police in some areas might counteract the benefits of this, as it made PWID reluctant to get sterile syringes in public settings. This risk is real, given Vietnamese media having broadcast cases of drug users getting caught while they went out to buy drugs during the lockdown [44]. Moreover, as outreach workers indicated, it is important to consider when financial difficulties in buying heroin during the lockdown and social distancing period might facilitate group use of heroin, and with it, potential equipment sharing.

The common practice of group meth use and smoking equipment sharing is worrisome. This norm related

to meth use has been reported in other studies among female sex workers, methadone patients, and people who use meth in general [22, 45, 46]. This practice carries with it a high HCV infection risk, as the HCV prevalence among PWID was nearly 90% [47], with low uptake of HCV care services in general [29]. Meth use could interfere with HIV treatment adherence and viral suppression, thereby increasing the likelihood of HIV transmission from people who use drugs to their sexual partners [48, 49]. Moreover, meth use facilitates tuberculosis transmission, a main cause of death among HIV-positive people who inject drugs [50]. Given that one reason for participants to use meth collectively was to cope with the boredom and stress related to the COVID-19 pandemic, it is likely that this practice will persist even after lockdown is lifted, since unemployment will likely continue to be an issue during the global recession [51].

The COVID-19 pandemic has created new challenges for sex workers in different parts of the world, especially those with untreated drug use disorders [9, 13, 52]. Our study shows that the financial pressure resulting from the containment measures could lead Vietnamese female sex workers to accept unsafe sex and risk their health. The link between drug use and acceptance of unsafe sex for more money has also been reported in a study among sex workers in Canada [53]. The livelihood of female sex workers during COVID-19 lockdowns is worrisome, given many had low education, no vocational training and no alternative jobs [54]. Moreover, compared to men, women who use drugs are more stigmatized and receive much less support from family and social services [55, 56]. The global crisis is exacerbating their vulnerabilities, and calls for action to be taken in order to minimize its devastating impact on this population.

Recommendations

Our findings suggest harm reduction programs for PWID in provinces with high service coverage, like Haiphong, could consider focusing on meth use and on female sex workers during the COVID-19 epidemic. Regarding meth use, education and information about the risk related to group meth use and distribution of smoking tools would minimize HCV transmission risk. However, further efforts should be made to bring female sex workers into methadone treatment to help them deal with their financial stress. Harm reduction programs could be integrated into current social aid programs targeting populations of low socioeconomic status, in order to provide them with a comprehensive service package [57]. Outreach work should receive further investment, as outreach workers not only represent the bridge between people in need and treatment services, but also bring material support to those most in need [52]. Future studies should continue

to document the living experiences of PWID and their risks as the epidemic and prolonged lockdown continue to affect most provinces in Vietnam.

Limitations

The study would reflect more truthfully the experiences of participants if it had been conducted during the lockdown rather than after the lockdown had finished. The COVID-19 mitigation measures also made ethnographical work—a more appropriate method for the study question—unfeasible. While we intentionally over-recruited women, the number of female sex workers is small. Moreover, we were not able to recruit venue-based women who worked under the management of pimps. These participants might be living in different risk environments than our street-based participants. Finally, the COVID-19 situation in Vietnam has become much more severe than it was in 2020. PWID might currently experience greater challenges than our report shows.

Conclusion

The COVID-19 pandemic disproportionately affected PWID and introduced new positive and negative influences on HIV/HCV risk behaviors. Most participants suffered from the financial impact of the epidemic mitigation measures. Unsafe injection might no longer be an issue among PWID in Haiphong, thanks to a combination of different factors including high awareness of injection-related HIV/HCV risk, and the availability of methadone treatment. However, group meth use and smoking equipment sharing as a common practice among PWID might elevate the risk for HCV, HIV and COVID-19 infection. Meth use as a means to cope with boredom and the stress related to COVID-19 might continue beyond the lockdown period. Female sex workers, especially those who were active heroin users, suffered most from COVID-related financial pressure and might resort to unsafe sex. These insights could orient harm reduction programs during this global crisis.

Abbreviations

DRIVE: DRug use and Infections in ViEtnam; FGD: Focus group discussion; PWID: People who inject drugs.

Acknowledgements

We thank our participants for sharing their insights with us. We are grateful for our peer outreach worker colleagues for helping with this work.

Authors' contributions

NTT designed the qualitative component, conducted qualitative data collection and analysis and led the writing of this article. HTG acquired funding for the study and supervised data collection and analysis. NQD analyzed quantitative data. NHA, LAN conducted data collection and preliminary analysis. DL, NN, DDJ, DTH, NNTT, KTHO, PMK, LSM, LM designed the study and contributed to the interpretation of the data. DR acquired funding for the study and supervised data collection and analysis. LMG designed the qualitative component

and supervised qualitative data collection and analysis. All authors read and approved the final manuscript.

Funding

The study received funding from ANRS-MIE (French research agency on HIV/AIDS, viral hepatitis and emerging infectious diseases) (ANRS COV22 DRIVE-COVID). ANRS-MIE was not involved in the design of the study, and collection, analysis, and interpretation of data, or in writing the manuscript.

Availability of data and materials

The datasets generated during and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Haiphong University of Medicine and Pharmacy Ethics Committee approved this study.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Centre for Training and Research on Substance Use and HIV, Hanoi Medical University, Room 211B, Building E3, #1 Ton That Tung Street, Hanoi, Vietnam. ²Hai Phong University of Medicine and Pharmacy, Haiphong, Vietnam. ³Pathogenesis and Control of Chronic and Emerging Infections, INSERM, Etablissement Français du Sang, University of Montpellier, Montpellier, France. ⁴Infectious Diseases Department, Caremeau University Hospital, Nîmes, France. ⁵New-York University College of Public Health, New York, NY, USA. ⁶Supporting Community Development Initiatives, Hanoi, Vietnam. ⁷CESP Inserm UMRS 1018, Paris Saclay University, Pierre Nicole Center, French Red Cross, 27 rue Pierre Nicole, 75005 Paris, France.

Received: 22 September 2021 Accepted: 15 December 2021

Published online: 29 January 2022

References

- Nanda S. Inequalities and COVID-19. In: Nanda S, editor. COVID-19: two volume set. Abingdon: Taylor & Francis; 2021.
- Dubey MJ, Ghosh R, Chatterjee S, Biswas P, Chatterjee S, Dubey S. COVID-19 and addiction. *Diabetes Metab Syndr*. 2020;14(5):817–23.
- Grebely J, Cerdá M, Rhodes T. COVID-19 and the health of people who use drugs: What is and what could be? *Int J Drug Policy*. 2020;83:102958.
- Jacka BP, Phipps E, Marshall BDL. Drug use during a pandemic: convergent risk of novel coronavirus and invasive bacterial and viral infections among people who use drugs. *Int J Drug Policy*. 2020;83:102895.
- Jemberie WB, Stewart Williams J, Eriksson M, Grönlund A-S, Ng N, Blom Nilsson M, et al. Substance use disorders and COVID-19: multi-faceted problems which require multi-pronged solutions. *Front Psychiatry*. 2020;11:714.
- Mota P. Avoiding a new epidemic during a pandemic: the importance of assessing the risk of substance use disorders in the COVID-19 era. *Psychiatry Res*. 2020;290:113142.
- Carrico AW, Horvath KJ, Grov C, Moskowitz JT, Pahwa S, Pallikuth S, et al. Double jeopardy: methamphetamine use and HIV as risk factors for COVID-19. *AIDS Behav*. 2020;24(11):3020–3.
- Wei Y, Shah R. Substance use disorder in the COVID-19 pandemic: a systematic review of vulnerabilities and complications. *Pharmaceuticals*. 2020;13(7):155.
- Platt L, Elmes J, Stevenson L, Holt V, Rolles S, Stuart R. Sex workers must not be forgotten in the COVID-19 response. *The Lancet*. 2020;396(10243):9–11.
- Dietze PM, Peacock A. Illicit drug use and harms in Australia in the context of COVID-19 and associated restrictions: anticipated consequences and initial responses. *Drug Alcohol Rev*. 2020;39(4):297–300.
- Karimi-Sari H, Sharafi H, Rezaee-Zavareh MS, Alavian SM. Harm reduction during the COVID-19 outbreak in Iran. *Lancet Psychiatry*. 2020;7(8):e57.
- Schlosser A, Harris S. Care during COVID-19: drug use, harm reduction, and intimacy during a global pandemic. *Int J Drug Policy*. 2020;83:102896.
- Callander D, Meunier É, DeVeau R, Grov C, Donovan B, Minichiello V, et al. Investigating the effects of COVID-19 on global male sex work populations: a longitudinal study of digital data. *Sex Transm Infect*. 2021;97(2):93–8.
- Starks TJ, Jones SS, Sauermilch D, Benedict M, Adebayo T, Cain D, et al. Evaluating the impact of COVID-19: a cohort comparison study of drug use and risky sexual behavior among sexual minority men in the U.S.A. *Drug Alcohol Depend*. 2020;216:108260.
- Wongsamuth N. FEATURE-Concern over “chemsex” grows as COVID-19 shuts Thailand’s LGBT+ bars. *Thompson Reuters Foundation* [Internet]. 2021 May 18 [cited 2021 Nov 5]. <https://www.reuters.com/article/thailand-lgbt-drugs-idUSL8N2BP44C>.
- Vo HS. Updates on the HIV epidemic in Vietnam. Conference: HIV and substance use in Vietnam—25 years of research and training; 2019. Hanoi, Vietnam.
- Hien NT, Long NT, Huan TQ. HIV/AIDS epidemics in Vietnam: evolution and responses. *AIDS Educ Prev*. 2004;16(supplement_a):137–54.
- Edington C, Bayer R. When grammars collide: Harm reduction, drug detention and the challenges of international policy reform efforts in Vietnam. *Glob Public Health*. 2013;8(SUPPL.1):S75–91.
- HIV/AIDS Data Hub for the Asia Pacific. HIV and AIDS in Vietnam [Internet]. 2021 [cited 2021 March 24]. <https://www.aidsdatahub.org>.
- Molès J-P, Vallo R, Khue PM, Huong DT, Oanh KTH, Thoa NT, et al. HIV control programs reduce HIV incidence but not HCV incidence among people who inject drugs in HaiPhong, Vietnam. *Sci Rep*. 2020;10(1):6999.
- Michel L, Des Jarlais DC, Thi HD, Hai OK, Minh KP, Peries M, et al. Intravenous heroin use in Haiphong, Vietnam: need for comprehensive care including methamphetamine use-related interventions. *Drug Alcohol Depend*. 2017;179:198–204.
- Ho HT, Le GM, Dinh TT. Female sex workers who use amphetamine-type stimulants (ATS) in three cities of Vietnam: use and sexual risks related to HIV/AIDS. *Glob Public Health*. 2013;8(5):552–69.
- Berto A, Day J, Van Vinh CN, Thwaites GE, My NN, Baker S, et al. Current challenges and possible solutions to improve access to care and treatment for hepatitis C infection in Vietnam: a systematic review. *BMC Infect Dis*. 2017;17(1):260.
- Pollack, Todd M., Thwaites GE, Rabaa M, Choisy M, van Doorn R, Tan LV, et al. Emerging COVID-19 success story: Vietnam’s commitment to containment [Internet]. *Our World in Data*. 2021 [cited 2021 Aug 16]. <https://ourworldindata.org/covid-exemplar-vietnam>.
- Ministry of Health. COVID-19 updates [Internet]. [cited 2021 Sep 15]. <https://covid19.gov.vn/big-story/ca-nhat-dien-bien-dich-covid-19-171210901110957364.htm>.
- Nguyen TTM, Nguyen LT, Pham MD, Vu HH, Mulvey KP. Methadone maintenance therapy in Vietnam: an overview and scaling-up plan. *Adv Prev Med*. 2012;2012:732484.
- Des Jarlais DC, Huong DT, Oanh KT, Pham MK, Giang HT, Thanh NT, et al. Prospects for ending the HIV epidemic among persons who inject drugs in Haiphong, Vietnam. *Int J Drug Policy*. 2016;32:50–6.
- Des Jarlais DC, Huong DT, Oanh KTH, Feelemyer JP, Arasteh K, Khue PM, et al. Ending an HIV epidemic among persons who inject drugs in a middle-income country: extremely low HIV incidence among persons who inject drugs in Hai Phong, Viet Nam. *AIDS*. 2020;34(15):2305–11.
- Thinh VT, Phuong DT, Hoa VD, Giang LM. Reported low uptake of HCV testing among people who inject drugs in urban Vietnam. *Biomed Res Int*. 2020;2020:e3701379.
- Rhodes T. The ‘risk environment’: a framework for understanding and reducing drug-related harm. *Int J Drug Policy*. 2002;13(2):85–94.
- Moore D, Fraser S. Putting at risk what we know: reflecting on the drug-using subject in harm reduction and its political implications. *Soc Sci Med*. 2006;62(12):3035–47.
- Rhodes T. Risk environments and drug harms: a social science for harm reduction approach. *Int J Drug Policy*. 2009;20(3):193–201.

33. Ayano G, Tulu M, Haile K, Assefa D, Habtamu Y, Araya G, et al. A systematic review and meta-analysis of gender difference in epidemiology of HIV, hepatitis B, and hepatitis C infections in people with severe mental illness. *Ann Gen Psychiatry*. 2018;17(1):16.
34. Bajis S, Dore GJ, Hajarizadeh B, Cunningham EB, Maher L, Grebely J. Interventions to enhance testing, linkage to care and treatment uptake for hepatitis C virus infection among people who inject drugs: a systematic review. *Int J Drug Policy*. 2017;47:34–46.
35. Karki P, Shrestha R, Huedo-Medina TB, Copenhaver M. The impact of methadone maintenance treatment on HIV risk behaviors among high-risk injection drug users: a systematic review. *Evid Based Med Public Health*. 2016;2:e1229.
36. ATLAS.ti: The qualitative data analysis & research software [Internet]. ATLAS.ti. [cited 2021 Jun 7]. <https://atlasti.com/>.
37. Charmaz K. *Constructing grounded theory*. New York: SAGE; 2014. p. 417.
38. DRIVE-COVID study team. Evaluating the impact of lockdown measures to control the COVID-19 epidemic on HIV and HCV risk behaviours and access to prevention and care for people who inject drugs in Haiphong, Vietnam. 2021.
39. Bonn M, Palayew A, Bartlett S, Brothers TD, Tounesnard N, Tyndall M. Addressing the syndemic of HIV, hepatitis C, overdose, and COVID-19 among people who use drugs: the potential roles for decriminalization and safe supply. *J Stud Alcohol Drugs*. 2020;81(5):556–60.
40. Larney S, Peacock A, Leung J, Colledge S, Hickman M, Vickerman P, et al. Global, regional, and country-level coverage of interventions to prevent and manage HIV and hepatitis C among people who inject drugs: a systematic review. *Lancet Glob Health*. 2017;5(12):e1208–20.
41. O'Carroll A, Duffin T, Collins J. Harm reduction in the time of COVID-19: case study of homelessness and drug use in Dublin, Ireland. *Int J Drug Policy*. 2020;87:102966.
42. Perri M, Dosani N, Hwang SW. COVID-19 and people experiencing homelessness: challenges and mitigation strategies. *CMAJ*. 2020;192(26):E716–9.
43. Boeri M, Gardner M, Gerken E, Ross M, Wheeler J. "I don't know what fun is": examining the intersection of social capital, social networks, and social recovery. *Drugs Alcohol Today*. 2016;16(1):95–105.
44. Dang H. [Going out to buy drugs during lockdown, many addicts are arrested]. Ho Chi Minh City Public Security [Internet]. 2021 Aug 2 [cited 2021 Aug 19]. https://congan.com.vn/vu-an/len-di-mua-ma-tuy-nhieuk-ke-nghien-bi-bat_117433.html.
45. Hanoi Medical University C for R and T on SA and H. Harm reduction for key populations who use methamphetamine in Vietnam. An acceptability and feasibility evaluation report [Internet]. 2021 Jan [cited 2021 Jul 22]. https://blogbird.b-cdn.net/mainline-eng/Evaluation_HR_for_key_populations_who_use_meth_in_Vietnam1.pdf.
46. Thuy DTT, Trang NT, Hoa NH, Hoe HD, Lai NL, Anh VM, et al. Tailoring evidence-based interventions for methadone patients who use methamphetamine in a lower middle-income country. 2021.
47. Zhang L, Celentano DD, Le Minh N, Latkin CA, Mehta SH, Frangakis C, et al. Prevalence and correlates of HCV mono-infection, HIV and HCV coinfection among persons who inject drugs in Vietnam. *Eur J Gastroenterol Hepatol*. 2015;27(5):550–6.
48. Moore DJ, Blackstone K, Woods SP, Ellis RJ, Atkinson JH, Heaton RK, et al. Methamphetamine use and neuropsychiatric factors are associated with antiretroviral non-adherence. *AIDS Care*. 2012;24(12):1504–13.
49. Jin H, Ogunbajo A, Mimiaga MJ, Duncan DT, Boyer E, Chai P, et al. Over the influence: the HIV care continuum among methamphetamine-using men who have sex with men. *Drug Alcohol Depend*. 2018;192:125–8.
50. Vinh VH, Vallo R, Giang HT, Huong DT, Oanh KTH, Khue PM, et al. A cohort study revealed high mortality among people who inject drugs in Hai Phong, Vietnam. *J Clin Epidemiol*. 2021;139:38–48.
51. Chodorow-Reich G, Coglianese J. Projecting unemployment durations: A factor-flows simulation approach with application to the COVID-19 recession. *J Public Econ*. 2021;197:104398.
52. Kimani J, Adhiambo J, Kasiba R, Mwangi P, Were V, Mathenge J, et al. The effects of COVID-19 on the health and socio-economic security of sex workers in Nairobi, Kenya: emerging intersections with HIV. *Glob Public Health*. 2020;15(7):1073–82.
53. Deering KN, Lyons T, Feng CX, Nosyk B, Strathdee SA, Montaner JS, et al. Client demands for unsafe sex: the socio-economic risk environment for HIV among street and off-street sex workers. *J Acquir Immune Defic Syndr*. 2013;63(4):522–31.
54. Meena Saraswathi S, Aarathi P, Laxmi M. Locked down: sex workers and their livelihoods. *Econ Political Wkly*. 2015;7–8.
55. Haritavorn N. Surviving in two worlds: social and structural violence of Thai female injecting drug users. *Int J Drug Policy*. 2014;25(1):116–23.
56. Spooner C, Saktiawati AMI, Lazuardi E, Worth H, Subronto YW, Padmawati RS. Impacts of stigma on HIV risk for women who inject drugs in Java: a qualitative study. *Int J Drug Policy*. 2015;26(12):1244–50.
57. Government of Vietnam. Resolution 86/NQ-CP on urgent measures to prevent and control the Covid-19 epidemic. 2021.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

